arXiv:math/0703554v3 [math.CO] 22 Nov 2007

Graphs with many r-cliques have large complete r-partite subgraphs

Vladimir Nikiforov

Department of Mathematical Sciences, University of Memphis, Memphis TN 38152

February 2, 2008

Abstract

Let $r\geq 2$ and $c>0$. Every graph on n vertices with at least cn^{r} cliques on r vertices contains a complete r-partite subgraph with $r-1$ parts of size $\lfloor c^{r}\log n\rfloor$ and one part of size greater than $n^{1-c^{r-1}}$. This result implies the Erdős-Stone-Bollobás theorem, the essential quantitative form of the Erdős-Stone theorem.

Keywords: *clique; number of cliques; r-partite graph; Erdős-Stone theorem*

Main results

This note is part of an ongoing project aiming to renovate some classical results in extremal graph theory, see, e.g., [5], [12, 15].

Recall a result of Erdős and Stone [7]: every graph with n vertices and $\lceil cn^{2}\rceil$ edges contains a complete bipartite subgraph with each part of size $\lfloor a\log n\rfloor$, where $c,a>0$ are independent of n.

One generalization of this result stems from the fundamental theorem of Erdős and Stone [7]: every graph with n vertices and $\lceil(1-1/r+c)\,n^{2}/2\rceil$ edges contains a complete $(r+1)$-partite graph with each part of size $g\left(r,c,n\right)$, where, for r and c fixed, $g\left(r,c,n\right)$ tends to infinity with n. In [3] Bollobás and Erdős found that $g\left(r,c,n\right)=\Theta\left(\log n\right)$, and in [2], [4], [6], and [8] the function $g\left(r,c,n\right)$ was determined with great precision.

In this note we deduce the Erdős-Stone-Bollobás result from weaker premises. Also, in our setup c may be a function of n, say $c=1/\ln\ln n$ and the bounds on $g\left(r,c,n\right)$ remain meaningful; such results are beyond the scope of the papers mentioned above.

Our notation follows [1]. Thus, $K_{r}\left(s_{1},\ldots,s_{r}\right)$ denotes a complete r-partite graph with parts of size $s_{1},\ldots,s_{r}$, and an r-clique is a complete subgraph on r vertices. We write $E\left(G\right)$ for the edge set of a graph G and $K_{r}\left(G\right)$ for the set of its r-cliques; we let $e\left(G\right)=\left|E\left(G\right)\right|$ and $k_{r}\left(G\right)=\left|K_{r}\left(G\right)\right|$.

Given $M\subset K_{r}\left(G\right)$, we write $K_{s}\left(M\right)$ for the set of s-cliques contained in members of M. Given a subgraph $H\subset G$ such that $H=K_{r}\left(s_{1},\ldots,s_{r}\right)$, we say that M *covers* H if $E\left(H\right)\subset K_{2}\left(M\right)$ and H contains $\min\left\{s_{1},\ldots,s_{r}\right\}$ disjoint members of M.

Here is our main result.

Theorem 1 *Let $r\geq 2$, $c^{r}\ln n\geq 1$, and G be a graph with n vertices. Every set of at least cn^{r} r-cliques of G covers a $K_{r}\left(s,\ldots s,t\right)$ with $s=\lfloor c^{r}\ln n\rfloor$ and $t>n^{1-c^{r-1}}$.*

Remarks

- Using random graphs, it is easy to see that most graphs on n vertices contain no complete bipartite subgraphs with both parts larger than $C \log n$, for some $C > 0$, independent of n. Hence, Theorem 1 is essentially best possible.
- Theorem 1 implies the Erdős-Stone-Bollobás theorem. Indeed, in [9] (see also [11], Problem 11.8) it is proved that if $k_s(G) > 0$, then

  $$\frac{(s+1)\, k_{s+1}(G)}{s k_s(G)} - \frac{n}{s} \geq \frac{s k_s(G)}{(s-1)\, k_{s-1}(G)} - \frac{n}{s-1}.$$

  Thus, if $e(G) \geq (1 - 1/r + c)\, n^2/2$, we find that $k_{r+1}(G) > (c/r^r)\, n^{r+1}$, and so G contains a $K_{r+1}(s, \ldots s, t)$ with

  $$s = \left\lfloor (c/r^r)^{r+1} \ln n \right\rfloor, \quad t > n^{1-(c/r^r)^r}.$$

  This is slightly stronger than the Erdős-Stone-Bollobás result and is comparable with those established in [4].
- The exponent $1 - c^{r-1}$ in Theorem 1 is not the best one, but is simple.

We deduce Theorem 1 from a routine lemma, proved here for convenience.

Lemma 2 *Let F be a bipartite graph with parts A and B. Let $|A| = m$, $|B| = n$, $r \geq 2$, $(\ln n)^{-1/r} \leq c < 1/2$, and $s = \lfloor c^r \ln n \rfloor$. If $s \leq (c/2)\, m + 1$ and $e(F) \geq cmn$, then F contains a $K_2(s,t)$ with parts $S \subset A$ and $T \subset B$ such that $|S| = s$ and $|T| = t > n^{1-c^{r-1}}$.*

Proof Let

$$t = \max\{x : \text{there exists } K_2(s,x) \subset F \text{ with part of size } s \text{ in } A\}.$$

For any $X \subset A$, write $d(X)$ for the number of vertices joined to all vertices of X. By definition, $d(X) \leq t$ for each $X \subset A$ with $|X| = s$; hence,

$$t\binom{m}{s} \geq \sum_{X \subset A, |X|=s} d(X) = \sum_{u \in B} \binom{d(u)}{s}. \tag{1}$$

Set

$$f(x) = \begin{cases} \binom{x}{s} & \text{if } x \geq s-1 \\ 0 & \text{if } x < s-1, \end{cases}$$

and note that $f(x)$ is a convex function. Therefore,

$$\sum_{u \in B} \binom{d(u)}{s} = \sum_{u \in B} f(d(u)) \geq n f\left(\frac{1}{n} \sum_{u \in B} d(u)\right) = n\binom{e(F)/n}{s} \geq n\binom{cm}{s}.$$

Combining this inequality with (1), and after rearranging, we find that

$$t \geq n\frac{cm\left(cm-1\right)\cdots\left(cm-s+1\right)}{m\left(m-1\right)\cdots\left(m-s+1\right)} > n\left(\frac{cm-s+1}{m}\right)^{s} \geq n\left(\frac{c}{2}\right)^{s}$$
$$\geq n\left(e^{\ln(c/2)}\right)^{c^{r}\ln n} = n^{1+c^{r}\ln(c/2)} = n^{1+2c^{r-1}(c/2)\ln(c/2)}.$$

Since $c/2 < 1/4 < 1/e$ and $x\ln x$ is decreasing for $0 < x < 1/e$, we see that

$$t > n^{1+2c^{r-1}(c/2)\ln c/2} > n^{1+c^{r-1}(1/2)\ln 1/4} > n^{1-c^{r-1}},$$

completing the proof. □

Proof of Theorem 1 Let $M \subset K_r\left(G\right)$ satisfy $\left|M\right| \geq cn^{r}$. To prove that M covers a $K_r\left(s,\ldots s,t\right)$ with $s = \left\lfloor c^{r}\ln n\right\rfloor$ and $t > n^{1-c^{r-1}}$, we use induction on r.

Suppose $r = 2$ and let A and B be two disjoint copies of $V\left(G\right)$. Define a bipartite graph F with parts A and B, joining $u \in A$ to $v \in B$ if $uv \in M$. Set $s = \left\lfloor c^{2}\ln n\right\rfloor$ and note that $s \leq \left(c/2\right)n + 1$. Since $e\left(F\right) = 2\left|M\right| > cn^{2}$, Lemma 2 implies that F contains a $K\left(s,t\right)$ with $t > n^{1-c}$. Hence M covers a $K_2\left(s,t\right)$, proving the assertion for $r = 2$. Assume the assertion true for $2 \leq r' < r$.

For a set $N \subset K_r\left(G\right)$ and a clique $R \in K_{r-1}\left(N\right)$ let $d_N\left(R\right)$ be the number of members of N containing R. We first show that there exists $L \subset M$ with $\left|L\right| > \left(c/2\right)n^{r}$ such that $d_L\left(R\right) > cn$ for all $R \in K_{r-1}\left(L\right)$. Indeed, set $L = M$ and apply the following procedure.

```
While there exists an $R \in K_{r-1}(L)$ with $d_L(R) \leq cn$ do
    Remove from $L$ all $r$-cliques containing $R$.
```

When this procedure stops, we have $d_L\left(R\right) > cn$ for all $R \in K_{r-1}\left(L\right)$ and also

$$\left|M\right| - \left|L\right| \leq cn\left|K_{r-1}\left(M\right)\right| \leq cn\binom{n}{r-1} < \frac{c}{\left(r-1\right)!}n^{r} \leq \frac{c}{2}n^{r},$$

implying that $\left|L\right| > \left(c/2\right)n^{r}$, as claimed.

Since $K_{r-1}\left(L\right) \subset K_{r-1}\left(G\right)$ and

$$\left|K_{r-1}\left(L\right)\right| \geq r\left|L\right|/n > r\left(c/2\right)n^{r}/n = \left(rc/2\right)n^{r-1},$$

the induction assumption implies that $K_{r-1}\left(L\right)$ covers an $H = K_{r-1}\left(m,\ldots,m\right)$ with $m = \left\lfloor \left(rc/2\right)^{r-1}\ln n\right\rfloor$.

For a subgraph $X \subset G$ and a vertex $v \in V\left(G\right)$, write $X + v$ for the subgraph of G induced by $V\left(X\right) \cup \{v\}$. Let A be a set of m disjoint $(r-1)$-cliques in H that are members of $K_{r-1}\left(L\right)$. Define a bipartite graph F with parts A and $B = V\left(G\right)$, joining $R \in A$ to $v \in B$ if $R + v \in L$. Set $s = \left\lfloor c^{r}\ln n\right\rfloor$. Since $d_L\left(R\right) > cn$ for all $R \in K_{r-1}\left(L\right)$, we have $e\left(F\right) > cmn$. Also, we find that

$$s \leq c^{r}\ln n < \left(c/2\right)\left(rc/2\right)^{r-1}\ln n < \left(c/2\right)\left\lfloor\left(rc/2\right)^{r-1}\ln n\right\rfloor + 1 = \left(c/2\right)m + 1.$$

Lemma 2 implies that there exists $K_2\left(s,t\right) \subset F$ with parts $S \subset A$ and $T \subset B$ such that $\left|S\right| = s$ and $\left|T\right| = t > n^{1-c^{r-1}}$. Let H^{*} be the subgraph of H induced by the union of the members of S; clearly, $H^{*} = K_{r-1}\left(s,\ldots,s\right)$. Since $R + v \in L$ for all $v \in T$ and $R \in K_{r-1}\left(H^{*}\right)$, we see that L covers a $K_r\left(s,\ldots,s,t\right)$, completing the induction step and the proof. □

Concluding remarks

Finally, a word about the project mentioned in the introduction: in this project we aim to give wide-range results that can be used further, adding more integrity to extremal graph theory.

Acknowledgement Thanks to Béla Bollobás, Sasha Kostochka, and Cecil Rousseau for helpful discussions.

References

[1] B. Bollobás, *Modern Graph Theory,* Graduate Texts in Mathematics, 184, Springer-Verlag, New York (1998).

[2] B. Bollobás, P. Erdős, On the structure of edge graphs, *Bull. London Math. Soc.* 5 (1973), 317-321.

[3] B. Bollobás, P. Erdős, M. Simonovits, On the structure of edge graphs II, *J. London Math. Soc.* (2) 12 (1976), 219–224.

[4] B. Bollobás, Y. Kohayakawa, An extension of the Erdős-Stone theorem, *Combinatorica* (3) 14 (1994), 279–286.

[5] B. Bollobás, V. Nikiforov, Joints in graphs, to appear in *Discrete Math.*

[6] V. Chvátal, E. Szemerédi, On the Erdős-Stone theorem. *J. London Math. Soc.* (2) 23 (1981), 207–214.

[7] P. Erdős, A.H. Stone, On the structure of linear graphs, *Bull. Amer. Math. Soc.* 52 (1946), 1087–1091.

[8] Y. Ishigami, Proof of a conjecture of Bollobás and Kohayakawa on the Erdős-Stone theorem, *J. Combin. Theory Ser. B* (2) 85 (2002), 222–254.

[9] N. Khadžiivanov, V. Nikiforov, The Nordhaus-Stewart-Moon-Moser inequality, (Russian), *Serdica* (4) 4 (1978), 344–350.

[10] T. Kövari, V. Sós, P. Turán, On a problem of K. Zarankiewicz, *Colloquium Math.* 3 (1954), 50–57.

[11] L. Lovász, *Combinatorial problems and exercises*, North-Holland Publishing Co., Amsterdam-New York (1979).

[12] V. Nikiforov, Turan's theorem inverted, submitted for publication. Preprint available at *http://arxiv.org/abs/0707.3394*

[13] V. Nikiforov, Stability for large forbidden graphs, submitted for publication. Preprint available at *http://arxiv.org/abs/0707.2563*

[14] V. Nikiforov, Graphs with many copies of a given subgraph, submitted for publication. Preprint available at *http://arxiv:0711.3493*

[15] V. Nikiforov, Complete r-partite subgraphs of dense r-graphs, submitted for publication. Preprint available at *http://arxiv.org/abs/0711.1185*